

## Extracellular Ascorbate: A Potential First-Line of Defence against Ozone

Jeremy D. Barnes<sup>1</sup>, Tom Lyons<sup>1</sup>, Joanna Maddison<sup>1</sup>, Enikő Turcsányi<sup>2</sup>, Youbin Zheng<sup>3\*</sup> and Matthias Plöchl<sup>4</sup>

<sup>1</sup>*Air Pollution Laboratory, Department of Agricultural and Environmental Science  
Ridley Building, The University, Newcastle Upon Tyne, NE1 7RU, UK*

<sup>2</sup>*Biological Research Centre, Hungarian Academy of Sciences, Institute of Plant Biology  
H-6701 Szeged, Temesvári krt. 62, P.O. Box 521, Hungary*

<sup>3</sup>*Center for Global Environmental Research (CGER),  
National Institute for Environmental Studies (NIES),  
16-2, Onogawa, Tsukuba, Ibaraki 305-0053, Japan*

<sup>4</sup>*Institute of Agricultural Engineering - Bornim (ATB)*

*Max-Eyth-Allee 100, D-14469 Potsdam-Bornim, Germany*

*E-mail: j.d.barnes@ncl.ac.uk, t.m.lyons@ncl.ac.uk, jo.maddison@ncl.ac.uk,  
enikot@mucclaus.szbk.u-szeged.hu, mploechl@atb-potsdam.de*

### Abstract

The mechanisms underlying the differential sensitivity of plants to the ubiquitous air pollutant ozone (O<sub>3</sub>) are far from fully understood. There is, however, a growing realisation that, following the uptake of the pollutant into the leaf interior, the first reactions take place in the aqueous matrix associated with the leaf cell walls (i.e. the leaf apoplast). This compartment forms the primary boundary between atmosphere and biosphere. The leaf apoplast is known to contain several antioxidants that react readily with O<sub>3</sub> (and/or its primary dissolution products) to yield ostensibly harmless compounds. There is therefore the possibility that significant amounts of O<sub>3</sub> are scavenged (i.e. detoxified) prior to reaching the primary target - the plasmalemma. If this is the case, antioxidants situated in the leaf apoplast may afford an important first-line of defence against O<sub>3</sub>. Herein, we focus on the role played by one of these compounds, ascorbate (vitamin C), in screening the plasmalemma from O<sub>3</sub>-induced oxidative insult.

**Key Words:** Antioxidants, Apoplast, Ascorbate, Cell wall, Detoxification, Modelling, ozone

### Introduction

The past century has witnessed a steady rise in tropospheric ozone (O<sub>3</sub>) concentrations in the Northern hemisphere (Volz and Kley, 1988; Marenco *et al.*, 1994), plus a sharp increase in the frequency and duration of potentially damaging photochemical episodes (Stockwell *et al.*, 1997). Indeed, there is unequivocal evidence that current ground-level concentrations of the O<sub>3</sub> are high enough to depress crop yields (Heck *et al.*, 1983; Fuhrer *et al.*, 1997), affect the composition and diversity of unmanaged ecosystems (Davison and Barnes, 1998) and contribute to localised declines in tree vitality in parts of Europe, North and Central America and the Far East (Chappelka and Chevone, 1992; Sandermann *et al.*, 1997; Izuta, 1998). The phytotoxicity of O<sub>3</sub> arises primarily as a result of the oxidative damage it causes to plasmalemma constituents (Heath, 1980, 1987, 1988). The pollutant is taken-up into the leaf interior, via the stomates (Kersteins and Lendzian, 1989), where it reacts with constituents of

*\*Present address: Department of Plant Agriculture, University of Guelph, Guelph Ontario, N1G 2W1, Canada  
E-mail: yzheng@uoguelph.ca*

the aqueous matrix associated with the cell wall (the apoplast) to yield a suite of reactive oxygen species which, in addition to O<sub>3</sub>, result in the oxidation of sensitive components of the plasmalemma, and subsequently the cytosol (Heath, 1980; Chameides, 1989; Moldau, 1998). In this sense, the oxidative stress induced by O<sub>3</sub> shares similarities with the initial events associated with other plant pathologies and the downstream consequences of O<sub>3</sub>-induced oxidative damage are well-documented i.e. visible leaf injury (i.e. localised cell death), stimulated rates of respiration, the suppression of photosynthesis, enhanced emissions of ethylene, premature leaf senescence and shifts in resource partitioning (Harris and Bailey-Serres, 1994; Kangasjärvi *et al.*, 1994; Schraudner *et al.*, 1997). These responses are manifested in decreased plant growth, depressed crop yields and reduced quality of harvested components (Runeckles and Chevone, 1992; Davison and Barnes, 1998; Barnes *et al.*, 1999a, b; Turcsányi *et al.*, 2000a). Since the plasmalemma is the principle site of attack, the interception and detoxification of O<sub>3</sub> (and/or its reactive products) by constituents of the leaf apoplast may play a crucial role in averting cellular damage (Heath, 1988; Kelly *et al.*, 1995; Dietz, 1997; Cross *et al.*, 1998; Lyons *et al.*, 1999a). In this contribution, we focus on evidence supporting a potentially important role for extracellular ascorbate (ASC) in mediating the tolerance of plants to O<sub>3</sub>.

### Ozone resistance

Herein, the term 'O<sub>3</sub> resistance' is defined as the "ability to maintain growth and remain free from injury in a polluted environment" (*sensu* Roose *et al.*, 1982). Resistance need not be complete and, based on the conceptual model proposed by Tingey and Andersen (1991), is envisaged to be mediated through effects on pollutant uptake (avoidance through changes in stomatal conductance) and/or changes in the tolerance of plant tissues following uptake (through effects on metabolism resulting in an increased capacity for the detoxification of the pollutant and its potentially damaging reaction products).


There is considerable variation in O<sub>3</sub> resistance within, as well as between, species (see Figure 1 which shows data for plantain [*Plantago major*]). This variation, which has been shown in many cases to be heritable, has been capitalised upon in several plant breeding programmes - directed selection resulting in the improved O<sub>3</sub> resistance of strains of alfalfa (*Medicago sativa* L.), sweet corn (*Zea mays* L.), snap bean (*Phaseolus vulgaris* L.), eastern white pine (*Pinus strobus* L.) and aspen (*Populus tremuloides* Michx.) (Howell *et al.*, 1971; Cameron, 1975; Mebrahtu *et al.*, 1990; Karnosky, 1991; Karnosky *et al.*, 1992).

It is generally concluded from such breeding experiments that O<sub>3</sub> resistance is a quantitative trait, governed by the additive effects of several genes (Sand, 1960; Howell *et al.*, 1971; DeVos *et al.*, 1982; Roose, 1991). However, Macnair (1991) has questioned this view, providing evidence that in many cases it is inevitable that air pollution resistance has been identified as a polygenic trait. Hence, it remains possible that O<sub>3</sub> resistance is controlled by a relatively small number of major genes and several modifiers. There is incontrovertible evidence that at least one of these genes is connected with the biosynthesis of L-ascorbate (vitamin C).


### Evidence supporting a role for ascorbate in mediating ozone resistance

Ascorbate is an abundant and powerful antioxidant which serves essential metabolic functions in both plants and animals (Loewus 1980, 1988; Smirnoff, 1996; Smirnoff and

Pallanca, 1996; Noctor and Foyer, 1998). The central position occupied by ASC in the metabolic control of reactive oxygen species is illustrated in Figure 2.



**Fig. 1** Ozone resistance in a range of geographically-distinct populations of *Plantago major*. Ozone resistance assessed in terms of the % change in relative growth rate induced by a two-week exposure to 70 ppb ozone for 7 h d<sup>-1</sup>. Data from Reiling and Davison (1992) (open bars) and Lyons *et al.* (1997) (closed bars).



**Fig. 2** Schematic representation of the central position occupied by reduced ascorbate (ASC), and its oxidised forms, monodehydroascorbate radical (MDHA) and dehydroascorbate (DHA), in the metabolism of reactive oxygen species: ozone (O<sub>3</sub>); superoxide radical (O<sub>2</sub><sup>-</sup>); hydroxyl radical (OH<sup>•</sup>); singlet oxygen (<sup>1</sup>O<sub>2</sub>); Hydrogen peroxide (H<sub>2</sub>O<sub>2</sub>). Enzyme and other metabolites involved are: superoxide dismutase (SOD); catalase (CAT); ascorbate peroxidase (APX); monodehydroascorbate radical reductase (MDHAR); dehydroascorbate reductase (DHAR); glutathione reductase (GR); reduced glutathione (GSH); oxidised glutathione (GSSG). Redrawn from Barnes *et al.* (1999b).

There are several strong lines of evidence linking leaf ASC content with O<sub>3</sub> resistance: (i) research with *Arabidopsis* mutants. Work performed by Conklin and co-workers (1996, 1997, 1999) has shown that O<sub>3</sub>-resistance co-segregates with the capacity to synthesize ASC in a range of *Arabidopsis thaliana* L. mutants derived from *vtc1* (formerly *soz1*); an ultrasensitive genotype that accumulates only 25% of the wild type leaf ASC concentration due to a deficiency in a key enzyme of the ASC biosynthetic pathway, GDP-mannose pyrophosphorylase. (ii) Correlative evidence. Endogenous levels of ASC correspond with variations in O<sub>3</sub> resistance in some cases (Lee *et al.*, 1984; Bilodeau and Chevrier, 1998), but not others (Menser, 1964; Ranieri *et al.*, 1999). (iii) Manipulation of leaf ASC content. Treatments employed to enhance leaf ASC content e.g. feeding ASC to roots (Mächler *et al.*, 1995) or spraying foliage (Freebairn, 1960; Freebairn and Taylor, 1960) have been shown to afford additional protection against O<sub>3</sub>. This effect is illustrated in Figure 3 using some of the authors' data to show that the extra protection observed in ASC-sprayed leaves cannot be explained through reduced rates of O<sub>3</sub> uptake (i.e. protection occurred in ASC-sprayed leaves of common plantain (*P. major* L.) in the absence of shifts in stomatal conductance).




Fig. 3 The effect of foliar ascorbate application on the response of (i) CO<sub>2</sub> assimilation rate (μmol m<sup>-2</sup> s<sup>-1</sup>) and (ii) stomatal conductance to water vapour (mmol m<sup>-2</sup> s<sup>-1</sup>) in leaves of *Plantago major* exposed to charcoal/Purafil®-filtered air (CFA, open symbols) or CFA plus 400 ppb ozone (closed symbols). Squares represent control plants sprayed with 10 mM NaHCO<sub>3</sub>, circles represent plants sprayed with 10 mM Na-ascorbate. Data represent means (*n* = 4) ± SE. Different letters denote significant differences (*P* < 0.05). Redrawn from Zheng *et al.* (2000).

In contrast, treatments which reduce leaf ASC content have been shown to enhance sensitivity to O<sub>3</sub> (Menser, 1964; Moldau *et al.*, 1998). More recently, some researchers have employed the recently-identified biosynthetic precursor of ASC, L-galactono-1, 4-lactone (L-GL), to enhance the ASC content of plant tissue. Bilodeau and Chevrier (1998), utilising an O<sub>3</sub> sensitive 'colourless-mutant' of *Euglena gracilis*, demonstrated that the addition of L-GL to the growth medium resulted in elevated levels of ASC in the cells, and increased their tolerance (in terms of cell viability) to O<sub>3</sub> exposure. In parallel experiments conducted by some of the authors (J. Maddison, T. Lyons & J. Barnes, unpublished), leaf ASC content was doubled in radish (*Raphanus sativus* L.) fed 50mM L-GL without effects on stomatal conductance. The treatment was found to afford complete protection against the negative effects of O<sub>3</sub> (in terms of visible injury and effects on plant relative growth rate).

## Importance of the sub-cellular localisation of ascorbate in the interception of ozone


The first reactions of ozone occur at the air/liquid interface (i.e. the leaf apoplast). Around 1-2% of the ASC found in leaves is located at this boundary. Hence, the ASC concentration in leaf cell walls ranges from 10 to 4000  $\mu\text{M}$  (Castillo and Greppin, 1988; Takahama and Oniki, 1992; Polle *et al.*, 1990, 1995; Luwe and Heber, 1995; Vanacker *et al.*, 1998a; Lyons *et al.*, 1999b; Ranieri *et al.*, 1999; Turcsányi *et al.*, 2000b). This raises the question of how much of the incoming  $\text{O}_3$  can be scavenged by this pool of ASC, and what the background rate of ASC consumption is (since ASC also participates as a co-factor in several enzyme-based reactions). *In vitro* studies on pure solutions have revealed that the ASC/ $\text{O}_3$  reaction proceeds rapidly (second-order reaction rate constant is between  $4.8 \times 10^7$  and  $6.0 \times 10^7 \text{ M}^{-1} \text{ s}^{-1}$  at physiologically-relevant pHs [Giamalva *et al.*, 1985; Kanofsky and Sima, 1995a]). Experiments involving the ozonation of isolated biological fluids, e.g. apoplastic washing fluid (Kanofsky and Sima, 1995b), respiratory tract lining fluid (Kelly *et al.*, 1995; Cross *et al.*, 1998) and blood plasma (Cross *et al.*, 1992; van der Vliet *et al.*, 1995) reveal that ASC is consumed at a rate dependent upon the concentration and the duration of exposure to the pollutant. Moreover, these studies have demonstrated that ASC is oxidised prior to reaction with lipid and protein constituents. These findings substantiate reports from *in vivo* studies that  $\text{O}_3$  exposure results in the depletion of extracellular ASC (Castillo and Greppin, 1988; Luwe *et al.*, 1993; Polle *et al.*, 1995; Luwe and Heber, 1995; Mudway *et al.*, 1999; Turcsányi *et al.*, 2000b). However, some researchers - Jakob and Heber (1998), in particular - are not yet convinced that *in vitro* findings are reciprocated *in vivo*. These authors have found that ASC does not prevent changes in the fluorescence of oxidation-sensitive dyes vacuum-infiltrated in to the apoplast of spinach (*Spinacia oleracea* L.) leaves, in contrast to experiments carried out in the test tube.

The product of ASC oxidation, dehydroascorbic acid (DHA) is reduced to regenerate ASC, or is rapidly and irreversibly hydrolysed to yield 2, 3-diketogulonic acid and an array of degradation products (Loewus, 1980, 1988; Smirnoff 1996; Deutsch, 1998a). Since DHA cannot be reduced efficiently in the apoplast, it is believed to be returned to the cytosol for recycling (Castillo and Greppin, 1984; Polle *et al.*, 1990; Luwe *et al.*, 1993). This view is supported by the presence of a carrier-mediated system on the plasma membrane for the transport of ASC/DHA (Rautenkranz *et al.*, 1994; Foyer and Lelandais, 1996; Horemans *et al.*, 1996). This system displays a higher affinity for DHA than for ASC (Horemans *et al.*, 1997, 1998). Recent reports by Vanacker and co-workers (Vanacker *et al.*, 1998a, b, 1999) suggest that there is the possibility, at least in some species, that DHA may be enzymatically reduced back to ASC in the apoplast - this finding awaits further investigation. It is also possible that monodehydroascorbate radical is formed as an intermediate during the reaction of ASC and  $\text{O}_3$ . If this is the case, plasmalemma-bound monodehydroascorbate radical educates may facilitate the rapid regeneration of apoplastic ASC *in situ* (Navas *et al.*, 1994; Asard *et al.*, 1995; Bérczi and Møller, 1998).

## Estimation of ozone detoxification in the leaf apoplast

Semi-quantitative estimates of the degree of protection afforded by apoplastic ASC were first provided by Chameides (1989). This author's insightful approach attempted to describe the uptake of  $\text{O}_3$  from the atmosphere to the mesophyll cell wall. The study indicated that the ASC/ $\text{O}_3$  reaction in the leaf apoplast could provide a major sink for the pollutant, based on a

first-order loss co-efficient for the reaction of between  $3,000\text{--}60,000\text{ s}^{-1}$ ; governed by the rate of the reaction and the concentration of apoplastic ASC (taken to be  $50\text{--}1000\text{ }\mu\text{M}$ ). More recently, a model (SODA; Simulating Ozone Detoxification in the Leaf Apoplast) which extends the one-dimensional approach adopted by Chameides (1989) to encompass the re-supply of ASC into the leaf apoplast under  $\text{O}_3$  exposure has been developed (Plöchl *et al.*, 2000). SODA estimates  $\text{O}_3$  uptake through the usual corollary with a series of resistances to diffusion first described by Gaastra (1959). Sub-cellular distribution of ASC is estimated from the diffusional movement of the neutral form, ascorbic acid (AA), and the ‘trapping’ of this compound in compartments of differing pH (*sensu* Slovik *et al.*, 1992). The detoxification of  $\text{O}_3$  in the leaf apoplast through direct reaction with ASC is modelled using a biomolecular reaction rate of  $4.8 \times 10^7\text{ M}^{-1}\text{ s}^{-1}$  (Kanofsky and Sima, 1995a). A schematic representation of the modelled resistance-reaction network is presented in Figure 4.



**Fig. 4** Schematic representation of the diffusion-reaction network modelled in SODA; considering the uptake of ozone ( $\text{O}_3$ ) from the air ( $\text{EXT}$ ) through the laminar boundary, stoma, intercellular air space ( $\text{INT}$ ), and its dissolution and reaction with ascorbate (ASC) in the aqueous phase of the cell wall (the apoplast,  $\text{APO}$ ), the distribution of ASC between cell compartments, as well as the transfer of dehydroascorbic acid (DHA) from apoplast to cytosol and its subsequent regeneration. Other abbreviations:  $c$  = concentration;  $\text{PL}$  = plasmalemma;  $\text{CHL}$  = chloroplast;  $\text{VAC}$  = vacuole. Redrawn from Plöchl *et al.* (2000).

The full complement of parameters necessary to rigorously test the theoretical prediction of mesophyll cell wall ASC concentrations, and  $\text{O}_3$  interception, have not yet been determined for an archetypal plant under standardised conditions. However, the best available dataset at this time, collected by Turcsányi *et al.* (2000b) for broad bean (*Vicia faba* L.) demonstrates a

reasonable agreement between modelled and measured ASC data (Figure 5). It is also interesting to note that recent work reveals (i) a positive correlation between the ASC content of the leaf apoplast and O<sub>3</sub> resistance (expressed in terms of reductions in growth rate induced by the pollutant) across a range of genotypes of *P. major* and *R. sativus* (Figure 6), and (ii) that pre-exposure of *P. vulgaris* plants to continuous darkness decreases the ASC content of the leaf apoplast by 80%, and results in an increase in injury to the plasmalemma upon exposure to acute O<sub>3</sub> (Moldau *et al.*, 1998).




Fig. 5 Relationship between measured and modelled concentration of ascorbate (mM) in the apoplast of *Vicia faba* leaves raised in CFA (charcoal/Purafil®-filtered air; squares) or ozone (O<sub>3</sub>) (CFA plus 75 nmol mol<sup>-1</sup> O<sub>3</sub> for 7 h d<sup>-1</sup>; circles) and exposed to CFA (open symbols) or 150 nmol mol<sup>-1</sup> O<sub>3</sub> (closed symbols). Line represents the least square regression fit;  $r^2 = 0.48$ ,  $P < 0.0001$ ,  $y = 0.885x + 0.099$  (replotted from Turcsányi *et al.*, 2000b).





Fig. 6 Relationship between ascorbate content of the leaf apoplast (nmol g<sup>-1</sup> FW) and ozone resistance in (i) populations of *Plantago major* ( $r^2 = 0.731$ ,  $P = 0.002$ ) and (ii) cultivars of *Raphanus sativus* ( $r^2 = 0.714$ ,  $P = 0.004$ ). Ozone resistance was assessed in terms of the % change in relative growth rate induced by a two-week exposure to 70 ppb ozone for 7 h d<sup>-1</sup>. Data represent means ( $n = 10-15$ )  $\pm$  SE (replotted from Barnes *et al.*, 1999b).

Model simulations emphasize the importance of pollutant concentrations in the atmosphere, stomatal conductance and the concentration of ASC in the leaf apoplast, in governing the flux of  $O_3$  that impinges on the plasmalemma (see Plöchl *et al.*, 2000). This point is emphasised in Figure 7 which shows (i) the effect of apoplastic ascorbate concentration on the flux of impinging on the plasmalemma. It is worthy of note that, at a concentration of ascorbate in the leaf apoplast of 500  $\mu M$ , ~60 % of the incoming  $O_3$  could be detoxified before it reaches the plasmalemma; (ii) the strong effect exerted by cell wall thickness on the detoxification of  $O_3$ , because of the way in which it modifies the length of the aqueous diffusion pathway to the plasmalemma (Nobel, 1991) and therefore the residence time of the pollutant in the cell wall. The thicker the wall, the greater the interception of the pollutant - reinforcing the conclusion of Chameides (1989); and (iii) the way in which the sub-cellular distribution of ASC is potentially affected by apoplast pH. The distribution of ASC between cell compartments is considered to be driven by a combination of pH and carrier-mediated transport systems, so shifts in the pH of the apoplast would be predicted to result in considerable changes in the concentration of ASC in the cell wall, and hence,  $O_3$  flux to the plasmalemma. This infers that environmental factors which alter apoplast pH e.g. water deficit (Hartung *et al.*, 1988), light (Mühling *et al.*, 1995), nitrogen supply (Hoffmann *et al.*, 1992) and gaseous pollutants including  $O_3$  itself (Heath, 1988; Wellburn, 1990; Pfanz and Oppmann, 1991; Moldau, 1998) may affect resistance to the pollutant in a predictable manner. This hypothesis is currently under investigation in the principle authors' laboratory.



**Fig. 7** The influence of (i) apoplastic ascorbate concentration ( $\mu M$ ), (ii) cell wall thickness ( $\mu m$ ) and (iii) apoplast pH on the simulated flux of ozone to the plasmalemma over a range of external ozone concentrations. Model inputs (unless otherwise shown) were: leaf ascorbate concentration = 6 mM; stomatal conductance to water vapour =  $200 \text{ mmol m}^{-2} \text{ s}^{-1}$ ; cell wall thickness =  $0.2 \mu m$ ; apoplast pH = 5.7; ascorbate/ozone reaction rate constant =  $4.8 \times 10^7 \text{ M}^{-1} \text{ s}^{-1}$ .

Present model formulations consider that  $O_3$  detoxification occurs solely the direct reaction of the pollutant with apoplastic ASC. There are numerous other apoplastic constituents that can scavenge  $O_3$  and its reactive products e.g. metabolites, such as polyamines and phenolic compounds (Bors *et al.*, 1989; Langebartels *et al.*, 1991; Eckey-Kaltenbach *et al.*, 1993, 1994), and enzymes, such as peroxidases and superoxide dismutase (Castillo *et al.*, 1984; Castillo and Greppin, 1988; Polle *et al.*, 1990; Streller and Wingsle, 1994; Ogawa *et al.*, 1996; Ranieri *et al.*, 1996; Vanacker *et al.*, 1998a, b, 1999; Lyons *et al.*, 1999b). Furthermore, the work of Deutsch (1998b) suggests that several of the oxidation products of ASC (including DHA) may act as strong antioxidants. There are also numerous reactions leading to the consumption of apoplastic ASC including the oxidative scission of cell wall polysaccharides *via* the formation of hydroxyl radicals (Fry, 1998), reaction with phenoxyl



radicals formed by cell wall peroxidases (Takahama and Oniki, 1992), regeneration of  $\alpha$ -tocopherol in the plasmalemma (Smirnoff, 1996) and the provision of substrate for ASC-dependant peroxidases (Castillo and Greppin, 1988) and oxidases (Takahama and Oniki, 1994). Our ultimate objective is to extend SODA to account for these reactions. However, a better understanding of the complex biochemistry of the apoplast is needed before this goal can be realized.

## Conclusions

Model simulations, allied to the available experimental data, indicate that extracellular ASC may play a significant role in the detoxification of  $O_3$  under environmentally-relevant conditions. Modulation of ASC content through various treatments leads to predictable shifts in  $O_3$  resistance; suggesting that it maybe possible to manipulate resistance *via* transgenic technology, particularly given recent advances in the characterisation of the biosynthetic pathway of ASC in plants. However, the data presented suggest that the protection against  $O_3$  afforded by apoplastic ASC is unlikely to complete. It appears likely that additional mechanisms may be important in capturing  $O_3$  in the leaf apoplast. Future work should be directed towards elucidating the nature of these extracellular defences and to the manipulation of these defences through present-day gene technology.

## References

- Asard, H., Horemans, N. and Caubergs, R.J. (1995) Involvement of ascorbic acid and a *b*-type cytochrome in plant plasma membrane redox reactions. *Protoplasma* 184: 36-41.
- Barnes, J.D., Balaguer, L., Manrique, E. and Davison, A.W. (1999a) Resistance to air pollutants: from cell to community. In: *Handbook of Functional Ecology*. Valladres, F. and Pugnaire, F.I. (eds.). 901pp. Marcel Dekker Inc., New York, 735-770.
- Barnes, J.D., Bender, J., Lyons, T. and Borland, A.M. (1999b) Natural and man-made selection for air pollution resistance. *Journal of Experimental Botany* 50: 1423-1435.
- Bérczi, A. and Møller, I.M. (1998) NADH-monodehydroascorbate oxidoreductase is one of the redox enzymes in spinach leaf plasma membranes. *Plant Physiology* 116: 1029-1036.
- Bilodeau, C. and Chevrier, N. (1998) Endogenous ascorbate level modulates ozone tolerance in *Euglena gracilis* cells. *Plant Physiology and Biochemistry* 36: 695-702.
- Bors, W., Langebartels, C., Michel, C. and Sandermann, H. (1989) Polyamines as protectants against ozone damage. *Phytochemistry* 28: 1589-1595.
- Cameron, J.W. (1975) Inheritance in sweet corn for resistance to acute ozone injury. *Journal of the American Society of Horticultural Science* 100: 577-579.
- Castillo, F.J. and Greppin, H. (1988) Extracellular ascorbic acid and enzyme activities related to ascorbic acid metabolism in *Sedum album* L. leaves after ozone exposure. *Environmental and Experimental Botany* 28: 231-238.
- Castillo, F.J., Penel, C. and Greppin, H. (1984) Peroxidase release induced by ozone in *Sedum album* leaves. Involvement of  $Ca^{2+}$ . *Plant Physiology* 74: 846-851.
- Chameides, W.L. (1989) The chemistry of ozone deposition to plant leaves: role of ascorbic acid. *Environmental Science and Technology* 23: 595-600.
- Chappelka, A.H. and Chevone, B.I. (1992) Tree responses to ozone. In: *Surface-Level Ozone Exposures and their Effects on Vegetation*. Lefohn, A.S. (ed.). 526pp. Lewis Publishers, Chelsea, 271-324.
- Conklin, P.L., Norris, S.R., Wheeler, G.L., Williams, E.H., Smirnoff, N. and Last, R.L. (1999) Genetic evidence for the role of GDP-mannose in plant ascorbic acid (vitamin C) biosynthesis. *Proceedings of the National Academy of Sciences, USA* 96: 4198-4203.

- Conklin, P.L., Pallanca, J.E., Last, R.L. and Smirnoff, N. (1997) L-Ascorbic acid metabolism in the ascorbate-deficient *Arabidopsis* mutant *vtc1*. *Plant Physiology* 115: 1277-1285.
- Conklin, P.L., Williams, E.H. and Last, R.L. (1996) Environmental stress sensitivity of an ascorbic acid-deficient *Arabidopsis* mutant. *Proceedings of the National Academy of Sciences, USA* 93: 9970-9974.
- Cross, C.E., Motchnik, P.A., Bruener, B.A., Jones, D.A., Kaur, H., Ames, B.N. and Halliwell, B. (1992) Oxidative damage to plasma constituents by ozone. *FEBS Letters* 298: 269-272.
- Cross, C.E., van der Vliet, A., Louie, S., Thiele, J.J., Halliwell, B. (1998) Oxidative stress and antioxidants at biosurfaces: plants, skin, and respiratory tract surfaces. *Environmental Health Perspectives* 106: 1241-1251.
- Davison, A.W. and Barnes, J.D. (1998) Impacts of ozone on wild plants. *New Phytologist* 139: 135-151.
- DeVos, N.E., Hill, R.R., Pell, E.J. and Cole, R.H. (1982) Quantitative inheritance of ozone resistance in potato. *Crop Science* 22: 992-995.
- Deutsch, J.C. (1998a) Ascorbic acid oxidation by hydrogen peroxide. *Analytical Biochemistry* 255: 1-7.
- Deutsch, J.C. (1998b) Oxygen-accepting antioxidants which arise during ascorbate oxidation. *Analytical Biochemistry* 265: 238-245.
- Dietz, K.-J. (1997) Functions and responses of the leaf apoplast under stress. *Progress in Botany* 58: 221-254.
- Eckey-Kaltenbach, H., Heller, W., Sonnenbichler, J., Zettl, I., Schäfer, W., Ernst, D. and Sandermann, H. (1993) Oxidative stress and plant secondary metabolism: 6''-O-malonylapin in parsley. *Phytochemistry* 34: 687-691.
- Eckey-Kaltenbach, H., Ernst, D., Heller, W. and Sandermann, H. (1994) Biochemical plant response to ozone. IV. Cross-induction of defensive pathways in parsley (*Petroselinum crispum* L.) plants. *Plant Physiology* 104: 67-74.
- Foyer, C.H. and Lelandais, M. (1996) A comparison of the relative rates of transport of ascorbate and glucose across the thylakoid, chloroplast and plasmalemma membranes of pea leaf mesophyll cells. *Journal of Plant Physiology* 148: 391-398.
- Freebairn, H.T. (1960) The prevention of air pollution damage to plants by use of vitamin C sprays. *Journal of the Air Pollution Control Association* 10: 314-317.
- Freebairn, H.T. and Taylor, O.C. (1960) Prevention of plant damage from airborne oxidising agents. *Proceedings of the American Society of Horticultural Science* 76: 693-699.
- Fry, S.C. (1998) Oxidative scission of plant cell wall polysaccharides by ascorbate-induced hydroxyl radicals. *Biochemical Journal* 332: 507-515.
- Fuhrer, J., Skärby, L. and Ashmore, M.R. (1997) Critical levels for ozone effects on vegetation in Europe. *Environmental Pollution* 97: 91-106.
- Gaastra, P. (1959) Photosynthesis of crop plants as influenced by light, carbon dioxide, temperature and stomatal diffusion resistance. *Mededelingen Landbouwhogeschool te Wageningen* 13: 1-68.
- Giamalva, D.H., Church, D.F. and Pryor, W.A. (1985) A comparison of the rates of ozonation of biological antioxidants and oleate and linoleate esters. *Biochemical and Biophysical Research Communications* 133: 773-779.
- Harris, M.J. and Bailey-Serres, J.N. (1994) Ozone effects on gene expression and molecular approaches to breeding for air pollution resistance. In: *Stress-Induced Gene Expression in Plants*. Basra, A.S. (ed.). 512pp. Harwood Academic, Singapore, 185-207.
- Hartung, W., Radin, J.W. and Hendrix, D.L. (1988) Absciscic acid movement into the apoplastic solution of water-stressed cotton leaves. *Plant Physiology* 86: 908-913.
- Heath, R.L. (1980) Initial events in injury to plants by air pollutants. *Annual Review of Plant Physiology* 31: 395-431.
- Heath, R.L. (1987) The biochemistry of ozone attack on the plasma membranes of plant cells. *Recent Advances in Phytochemistry* 21: 29-54.
- Heath, R.L. (1988) Biochemical mechanisms of pollutant stress. In: *Assessment of Crop Loss from Air Pollutants*. Heck, W.W., Taylor, O.C. and Tingey, D.T. (eds). 552pp. Elsevier Science, New York, 259-286.
- Heck, W.W., Adams, R.M., Cure, W.W., Heagle, A.S., Heggestad, H.E., Kohut, R.J., Kress, L.W., Rawlings, J.O. and Taylor, O.C. (1983) A reassessment of crop loss from ozone. *Environmental Science and Technology* 12: 572A-581A.
- Hoffmann, B., Plänker, R. and Mengal, K. (1992) Measurements of pH in the apoplast of sunflower leaves by means of fluorescence. *Physiologia Plantarum* 84: 146-153.
- Horemans, N., Asard, H. and Caubergs, R.J. (1996) Transport of ascorbate into plasma membrane vesicles of *Phaseolus vulgaris* L. *Protoplasma* 194: 177-185.

- Horemans, N., Asard, H. and Caubergs, R.J. (1997) The ascorbate carrier of higher plant plasma membranes preferentially translocates the fully oxidized (dehydroascorbate) molecule. *Plant Physiol* 114: 1247-1253.
- Horemans, N., Asard, H., Van Gestelen, P. and Caubergs, R.J. (1998) Facilitated diffusion drives transport of oxidised ascorbate molecules into purified plasma membrane vesicles of *Phaseolus vulgaris*. *Physiologica Plantarum* 104: 783-789.
- Howell, R.K., Devine, T.E. and Hanson, C.H. (1971) Resistance of selected alfalfa strains to ozone. *Crop Science* 11: 114-5.
- Izuta, T. (1998) Ecophysiological responses of Japanese forest tree species to ozone, simulated acid rain and soil acidification. *Journal of Plant Research* 111: 471-480.
- Jakob, A. and Heber, U. (1998) Apoplastic ascorbate does not prevent the oxidation of fluorescent amphiphilic dyes by ambient and elevated concentrations of ozone in leaves. *Plant and Cell Physiology* 39: 313-322.
- Kangasjärvi, J., Talvinen, J., Utriainen, M. and Karjalainen, R. (1994) Plant defence systems induced by ozone. *Plant, Cell and Environment* 17: 783-794.
- Kanofsky, J.R. and Sima, P.D. (1995a) Reactive absorption of ozone by aqueous biomolecule solutions: implications for the role of sulfhydryl compounds as targets for ozone. *Archives of Biochemistry and Biophysics* 316: 52-62.
- Kanofsky, J.R. and Sima, P.D. (1995b) Singlet oxygen generation from the reaction of ozone with plant leaves. *J Biol Chem* 270: 7850-7852.
- Karnosky, D.F. (1991) Ecological genetics and changes in atmospheric chemistry: the application of knowledge. In: *Ecological genetics and air pollution*. Taylor G.E., Pitelka, L.F. and Clegg, M.T. (eds.). 260pp. Springer-Verlag, New York Berlin London Tokyo, 321-336.
- Karnosky, D.F., Gagnon, Z.E., Reed, D.D. and Whitter, J.A. (1992) Growth and biomass allocation of symptomatic and asymptomatic *Populus tremuloides* clones in response to seasonal ozone exposures. *Canadian Journal of Forest Research* 22: 1785-1788.
- Kelly, F.J., Mudway, I., Krishna, M.T. and Holgate, S.T. (1995) The free radical basis of air pollution: focus on ozone. *Respiratory Medicine* 89: 647-656.
- Kersteins, G. and Lendzian, K.J. (1989) Interactions between ozone and plant cuticles. 1. Ozone deposition and permeability. *New Phytologist* 112: 13-19.
- Langebartels, C., Kerner, K., Leonardi, S., Schraudner, M., Trost, M., Heller, W. and Sandermann, H. (1991) Biochemical plant responses to ozone. I. Differential induction of polyamine and ethylene biosynthesis in tobacco. *Plant Physiology* 95: 882-889.
- Lee, E.H., Jersey, J.A., Gifford, C. and Bennett, J. (1984) Differential ozone tolerance in soybean and snap beans: analysis of ascorbic acid in O<sub>3</sub>-suseptible and O<sub>3</sub>-resistant cultivars by high-performance liquid chromatography. *Environmental and Experimental Botany* 24: 331-341.
- Loewus, F.A. (1980) L-ascorbic acid: metabolism, biosynthesis, function. In: *The Biochemistry of Plants, Volume 3*. Preiss, J. (ed.). 500pp. Academic Press, New York, 77-99.
- Loewus, F.A. (1988) Ascorbic acid and its metabolic products. In: *The Biochemistry of Plants, Volume 14*. Preiss, J. (ed.). 480pp. Academic Press, New York, 85-107.
- Luwe, M. and Heber, U. (1995) Ozone detoxification in the apoplasm and symplasm of spinach, bean and beech leaves at ambient and elevated concentrations of ozone in air. *Planta* 197: 448-455.
- Luwe, M.W.F., Takahama, U. and Heber, U. (1993) Role of ascorbate in detoxifying ozone in the apoplast of spinach (*Spinacia oleracea* L.) leaves. *Plant Physiology* 101: 969-976.
- Lyons, T.M., Barnes, J.D. and Davison, A.W. (1997) Relationships between ozone resistance and climate in European populations of *Plantago major*. *New Phytologist* 136: 503-510.
- Lyons, T., Plöchl, M., Turcsányi, E. and Barnes, J. (1999a) Extracellular antioxidants: a protective screen against ozone? In: *Environmental Pollution and Plant Responses*. Agrawal, S., Agrawal, M. and Krizek, D.T. (eds.). 393pp. CRC Press/Lewis Publishers, Boca Raton, 183-201.
- Lyons, T., Ollerenshaw, J.H. and Barnes, J.D. (1999b) Impacts of ozone on *Plantago major*: apoplastic and symplastic antioxidant status. *New Phytologist* 141: 253-263.
- Mächler, F., Wasescha, M.R., Krieg, F. and Oertli, J.J. (1995) Damage by ozone and protection by ascorbic acid in barley leaves. *Journal of Plant Physiology* 147: 469-473.
- Macnair, M.R. (1991) Genetics of the resistance of plants to pollutants. In: *Ecological Genetics and Air Pollution*. Taylor GE, Pitelka LF, Clegg MT, eds. 360pp. Springer-Verlag, New York Berlin London Tokyo, 127-136.
- Marengo, A., Gouget, H., Nedelec, P. and Pages, J.-P. (1994) Evidence of a long term increase in tropospheric ozone from Pic Du Midi data series: consequences - positive radiative forcing. *Journal of Geophysical Research* 99: 16617-16632.

- Mebrahtu, T., Mersie, W., Rangappa, M. (1990) Inheritance of ambient ozone insensitivity in common bean (*Phaseolus vulgaris* L.). *Environmental Pollution* 67: 79-89.
- Menser, H.A. (1964) Response of plants to air pollutants. III. A relation between ascorbic acid levels and ozone susceptibility of light-preconditioned tobacco plants. *Plant Physiology* 39: 564-567.
- Moldau, H. (1998) Hierarchy of ozone scavenging reaction in the plant cell wall. *Physiologia Plantarum* 104: 617-622.
- Moldau, H., Bichele, I. and Hüve, K. (1998) Dark-induced ascorbate deficiency in leaf cell walls increases plasmalemma injury under ozone. *Planta* 207: 60-66.
- Mudway, I.S., Krishna, M.T., Frew, A.J., MacLeod, D., Sandstrom, T., Holgate, S.T. and Kelly, F.J. (1999) Compromised concentrations of ascorbate in fluid lining the respiratory tract in human subjects after exposure to ozone. *Occupational and Environmental Medicine* 56: 473-481.
- Mühling, K.H., Plieth, C., Hansen, U.-P. and Sattelmacher, B. (1995) Apoplastic pH of intact leaves of *Vicia faba* as influenced by light. *Journal of Experimental Botany* 46: 377-382.
- Navas, P., Villalba, J.M. and Córdoba, F. (1994) Ascorbate function at the plasma membrane. *Biochimica et Biophysica Acta* 1197: 1-13.
- Nobel, P.S. (1991) *Physicochemical and Environmental Plant Physiology*. Academic Press, San Diego, 635pp.
- Noctor, G. and Foyer, C.H. (1998) Ascorbate and glutathione: keeping active oxygen under control. *Annual Review of Plant Physiology and Plant Molecular Biology* 49: 249-279.
- Ogawa, K., Kanematsu, S. and Asada, K. (1996) Intra- and extra-cellular localization of "cytosolic" CuZn-superoxide dismutase in spinach leaf and hypocotyl. *Plant and Cell Physiology* 37: 790-799.
- Pfanz, H. and Oppmann, B. (1991) The possible role of apoplastic peroxidases in detoxifying the air pollutant sulphur dioxide. In: *Biochemical, Molecular and Physiological Aspects of Plant Peroxidases*. Lobarzewski, J., Greppin, H., Penel, C. and Gaspar, T. (eds.). 600pp. University Marie Curie-Sklodowska and University of Geneva, Lublin and Geneva, 401-417.
- Plöchl, M., Lyons, T., Ollerenshaw, J.H. and Barnes, J.D. (2000) Simulating ozone detoxification in the leaf apoplast through the direct reaction with ascorbate. *Planta* 210: 454-467.
- Polle, A., Chakrabarti, K., Schurmann, W. and Rennenberg, H. (1990) Composition and properties of hydrogen peroxide decomposing systems in extracellular and total extracts from needles of Norway spruce (*Picea abies* L., Karst.). *Plant Physiology* 94: 312-319.
- Polle, A., Wieser, G. and Havranek, W.M. (1995) Quantification of ozone influx and apoplastic ascorbate content in needles of Norway spruce trees (*Picea abies* L., Karst.) at high altitude. *Plant, Cell and Environment* 18: 681-688.
- Ranieri, A., Castagna, A., Padu, E., Moldau H., Rahi, M. and Soldatini, G.F. (1999) The decay of O<sub>3</sub> through direct reaction with cell wall ascorbate is not sufficient to explain the different degrees of O<sub>3</sub>-sensitivity in two poplar clones. *Journal of Plant Physiology* 154: 250-255.
- Ranieri, A., D'Urso, G., Nali, C., Lorenzini, G. and Soldatini, G.F. (1996) Ozone stimulates apoplastic antioxidant systems in pumpkin leaves. *Physiologia Plantarum* 97: 381-387.
- Rautenkrantz, A.A.F., Li, L., Mächler, F., Martinoia, E. and Oertli, J.J. (1994) Transport of ascorbic and dehydroascorbic acids across protoplast and vacuole membranes isolated from barley (*Hordeum vulgare* L. cv Gerbel) leaves. *Plant Physiology* 106: 187-193.
- Reiling, K. and Davison, A.W. (1992) Spatial variation in ozone resistance of British populations of *Plantago major* L. *New Phytologist* 122: 699-708.
- Roose, M.L. (1991) Genetics of response to atmospheric pollutants. In: *Ecological Genetics and Air Pollution*. Taylor, G.E., Pitelka, L.F. and Clegg, M.T. (eds.). 360pp. Springer-Verlag, New York Berlin London Tokyo, 111- 126.
- Roose, M.L., Bradshaw, A.D. and Roberts, T.M. (1982) Evolution of resistance to gaseous air pollution. In: *Effects of Gaseous Air Pollutants in Agriculture and Horticulture*. Unsworth, M.H. and Ormrod, D.P. (eds.). 532pp. Butterworth Scientific, London, 379-409.
- Runeckles, V.C. and Chevone, B.I. (1992) Crop responses to ozone. In: *Surface-Level Ozone Exposures and their Effects on Vegetation*. Lefohn, A.S. (ed.). 526pp. Lewis Publishers, Chelsea, 189-270.
- Sand, S.A. (1960) Weather fleck in shade tobacco as problem of interaction between genes and environment. *Tobacco Science* 4: 137-146.
- Sandermann, H., Wellburn, A.R. and Heath, R.L. (1997) *Forest Decline and Ozone: A Comparison of Controlled Chamber and Field Experiments*, *Ecological Studies* 127. Springer-Verlag, Berlin, 400pp.
- Schraudner, M., Langebartels, C. and Sandermann, H. (1997) Changes in the biochemical status of plant cells induced by the environmental pollutant ozone. *Physiologia Plantarum* 100: 274-280.

- Slovik, S., Baier, M. and Hartung, W. (1992) Compartmental distribution and redistribution of acid in intact leaves. I. Mathematical formulation. *Planta* 187: 14-25.
- Smirnoff, N. (1996) The function and metabolism of ascorbic acid in plants. *Annals of Botany* 78: 661-669.
- Smirnoff, N. and Pallanca, J.E. (1996) Ascorbate metabolism in relation to oxidative stress. *Biochemical Society Transactions* 24: 472-478.
- Stockwell, W.R., Kramm, G., Scheel, H.-E., Mohnen, V.A. and Seiler, W. (1997) Ozone formation, destruction and exposure in Europe and the United States. In: *Forest Decline and Ozone: a Comparison of Controlled Chamber and Field Studies. Ecological Studies Series Vol. 127*. Sandermann, H., Wellburn, A.R. and Heath, R.L. (eds.) 400pp. Springer-Verlag, Heidelberg, 1-38.
- Streller, S. and Wingsle, G. (1994) *Pinus sylvestris* L. needles contain extracellular CuZn superoxide dismutase. *Planta* 192: 195-201.
- Takahama, U. and Oniki, T. (1992) Regulation of peroxidase-dependant oxidation of phenolics in the apoplast of spinach leaves by ascorbate. *Plant and Cell Physiology* 33: 379-387.
- Takahama, U. and Oniki, T. (1994) The association of ascorbate and ascorbate oxidase in the apoplast with IAA-enhanced elongation of epicotyls from *Vigna angularis*. *Plant and Cell Physiology* 35: 257-266.
- Tingey, D.T. and Andersen, C.P. (1991) The physiological basis of differential plant sensitivity to changes in atmospheric quality. In: *Ecological Genetics and Air Pollution*. Taylor, G.E., Pitelka, L.F. and Clegg, M.T. (eds.) 360pp. Springer-Verlag, New York Berlin London Tokyo, 209-35.
- Turcsányi, E., Cardoso-Vilhena, J., Maddison, J., Gillespie, C., Balaguer, L., Ollerenshaw, J.H. and Barnes, J.D. (2000a). Impacts of tropospheric ozone: past, present and likely future. In: *Trace Gas Emissions and Plants*. Singh, S.N. (ed.) Kluwer Academic Publishers, The Netherlands, in press.
- Turcsányi, E., Lyons, T., Plöchl, M. and Barnes, J. (2000b) Does ascorbate in the mesophyll cell walls form the first line of defence against ozone? 1. Testing the concept using broad bean (*Vicia faba* L.). *Journal of Experimental Botany* 51: 901-910.
- Vanacker, H., Carver, T.L.W. and Foyer, C.H. (1998a) Pathogen-induced changes in the antioxidant status of the apoplast in barley leaves. *Plant Physiology* 117: 1103-1114.
- Vanacker, H., Harbinson, J., Ruisch, J., Carver, T.L.W. and Foyer, C.H. (1998b) Antioxidant defences of the apoplast. *Protoplasma* 205: 129-140.
- Vanacker, H., Carver, T.L.W. and Foyer, C.H. (1999) Changes in apoplastic antioxidants induced by powdery mildew attack in oat genotypes with race non-specific resistance. *Planta* 208: 444-452.
- Van der Vliet, A., O'Neill, C.A., Eiserich, J.P. and Cross, C.E. (1995) Oxidative damage to extracellular fluids by ozone and possible protective effects of thiols. *Archives of Biochemistry and Biophysics* 321: 43-50.
- Volz, A. and Kley, D. (1988) Evaluation of the Montsouris series of ozone measurements made in the nineteenth century. *Nature* 332: 240-242.
- Wellburn, A.R. (1990) Why are atmospheric oxides of nitrogen usually phytotoxic and not alternative fertilizers? *New Phytologist* 115: 395-402.
- Zheng, Y., Lyons, T., Ollerenshaw, J.H., Barnes, J.D. (2000) Ascorbate in the leaf apoplast is a factor mediating ozone resistance in *Plantago major*. *Plant Physiology and Biochemistry* 38: 403-411.